

ON GLOBAL REPRESENTATIONS OF THE SOLUTIONS OF LINEAR DIFFERENTIAL EQUATIONS AS A PRODUCT OF EXPONENTIALS

J. WEI AND E. NORMAN

1. Introduction. In this paper we shall consider solutions of the equation

$$(1) \quad \frac{dU(t)}{dt} = A(t)U(t), \quad U(0) = I,$$

where A and U are linear operators, and I is the identity operator. Our results will be applicable when the operator $A(t)$ can be written as

$$A(t) = \sum_{i=1}^m a_i(t)X_i, \quad m \text{ finite,}$$

where the $a_i(t)$ are scalar functions of t , and the operators X_i are independent of t . It is further required that the Lie algebra \mathfrak{L} generated by the X_i under the commutator product $[X_i, X_j] = X_iX_j - X_jX_i$ be of finite dimension l . The above is, of course, always true if A (and U) are finite matrix operators.

In 1954, W. Magnus [4] proved that if X_1, X_2, \dots, X_l is a basis for \mathfrak{L} , then the solution of (1) can be expressed in the form $U(t) = \exp(\sum_{i=1}^l g_i(t)X_i)$. This representation of U holds, however, only in a neighborhood of the origin. It has been shown by J. Mariani and W. Magnus [3] that even in the case of 2×2 matrices a global version of Magnus' result cannot be obtained without severe restrictions on $A(t)$.

We will show that if U is a solution of (1), it can be represented in the form

$$(2) \quad U(t) = \prod_{i=1}^l \exp(g_i(t)X_i).$$

This representation is global for all solvable Lie algebras, and for any real 2×2 system of equations.

The form (2) derives its principal utility from the fact that insight into the properties of $U(t)$ can be gained through a knowledge of the spectral properties of the *individual* operators X_i . Since the X_i 's are

Received by the editors July 26, 1962 and, in revised form, January 17, 1963.

constant, and often have simple physical interpretations, one has a good chance of obtaining their spectral properties.

2. **Preliminaries.** In the sequel it will be convenient to refer to the independent variable as “time.”

Suppose the linear operator $A(t)$ can be expressed in the form

$$(3) \quad A(t) = \sum_{i=1}^m a_i(t)X_i, \quad m \text{ finite,}$$

where the $a_i(t)$ are scalar functions of time, and X_1, X_2, \dots, X_m are time-independent operators. If m is chosen as small as possible, the X_i 's will be linearly independent. We shall denote by R the associative algebra generated by X_1, X_2, \dots, X_m over the field of complex numbers; and by \mathfrak{L} the Lie algebra generated by X_1, \dots, X_m under the commutator product $[X_i, X_j] = X_iX_j - X_jX_i$, i.e., the Lie elements of R . R may be infinite-dimensional, but we shall assume that \mathfrak{L} is of finite dimension l .

We will require the following two lemmas.

(i) (BAKER-HAUSDORFF). *If, $x, y \in \mathfrak{L}$, then $e^x y e^{-x} \in \mathfrak{L}$, and is given by the explicit formula*

$$(4) \quad e^x y e^{-x} = y + [x, y] + [x, [x, y]]/2! + [x[x, [x, y]]]/3! + \dots$$

A proof is given in Magnus [4].

If we use the notation, adx , to represent the linear operator on \mathfrak{L} defined by

$$\begin{aligned} (adx)y &= [x, y] \\ (adx)^2 y &= [x, [x, y]], \text{ etc.} \end{aligned} \quad x, y \in \mathfrak{L},$$

then the Baker-Hausdorff formula (4) can be rewritten as

$$(5) \quad e^x y e^{-x} = (e^{adx})y.$$

(ii) *Let X_1, X_2, \dots, X_l be a basis for \mathfrak{L} with the multiplication table $[X_i, X_j] = \sum_{k=1}^l \gamma_{ij}^k X_k$, $i, j = 1, 2, \dots, l$. Then*

$$(6) \quad \left(\prod_{j=1}^r \exp(g_j X_j) \right) X_i \left(\prod_{j=r}^1 \exp(-g_j X_j) \right) = \sum_{k=1}^l \xi_{ki} X_k, \\ r = 1, 2, \dots, l,$$

where each $\xi_{ki} = \xi_{ki}(g_1, \dots, g_r)$ is an analytic function of g_1 to g_r .

PROOF. Repeated application of the previous lemma shows that the left-hand side of (6) is in \mathfrak{L} , and hence can be written as a linear

combination of X_1 to X_l as asserted. It remains to show that the ξ_k are analytic. It is sufficient to prove the result for $r=1$ since an analytic function of an analytic function is again analytic. For $r=1$ we have from (4)

$$\begin{aligned} \exp(g_1 X_1) X_i \exp(-g_1 X_1) &= \exp(g_1 \text{ad} X_1) X_i \\ &= X_i + \sum_{n=1}^{\infty} \frac{g_1^n}{n!} (\text{ad} X_1)^n X_i. \end{aligned}$$

Now

$$(7) \quad (\text{ad} X_1)^n X_i = \sum \gamma_{i_1}^{i_1} \gamma_{i_2}^{i_2} \cdots \gamma_{i_n}^{i_n} X_{i_n}, \quad i_1, \cdots, i_n = 1, \cdots, l.$$

Let M be the maximum of $|\gamma_{ij}^k|$, $i, j, k = 1, \cdots, l$. Estimating each $\gamma_{i_{s-1}}^{i_s}$ in (7) by M we obtain

$$\left| \sum \gamma_{i_1}^{i_1} \cdots \right| \leq (lM)^n.$$

Since $(lM)^n/n!$ is the general term of a convergent series the lemma is proved.

3. The local theorem.

THEOREM 1. *Let $A(t)$ be given by (3), and let the Lie algebra \mathfrak{L} generated by $A(t)$ be of finite dimension l . Then there exists a neighborhood of $t=0$ in which the solution of the equation*

$$(8) \quad \frac{dU}{dt} = A(t)U, \quad U(0) = I$$

may be expressed in the form

$$(9) \quad U(t) = \exp(g_1(t)X_1) \exp(g_2(t)X_2) \cdots \exp(g_l(t)X_l),$$

where the $g_i(t)$ are scalar functions of time. Moreover, the $g_i(t)$ satisfy a set of differential equations which depend only on the Lie algebra \mathfrak{L} , and the $a_i(t)$'s.

PROOF. The representation (9) is immediate from the Magnus representation via the second canonical coordinate system. We will, however, prove this fact again in the course of deriving the differential equations satisfied by the $g_i(t)$.

First note that we might just as well write $A(t) = \sum_{i=1}^l a_i(t)X_i$ instead of $A(t) = \sum_{i=1}^m a_i(t)X_i$ by simply setting $a_i(t) \equiv 0$ for $i > m$. Note also that at time $t=0$, $U(0) = I$ is in the form (9) with all $g_i(t) = 0$.

Now let U be of the form (9). Since

$$(10) \quad \begin{aligned} \frac{dU}{dt} &= \sum_{i=1}^l g'_i(t) \left(\prod_{j=1}^{i-1} \exp(g_j X_j) X_i \prod_{j=i}^l \exp(g_j X_j) \right) \\ AU &= \sum_{i=1}^l a_i(t) X_i \cdot U. \end{aligned}$$

We obtain upon substitution of (10) into (8), and post-multiplication by the inverse operator U^{-1}

$$(11) \quad \begin{aligned} \sum_{i=1}^l a_i(t) X_i &= \sum_{i=1}^l g'_i(t) \left(\prod_{j=1}^{i-1} \exp(g_j X_j) X_i \prod_{j=i-1}^1 \exp(-g_j X_j) \right) \\ &= \sum_{i=1}^l g'_i(t) \left(\prod_{j=1}^{i-1} \exp(g_j ad X_j) \right) X_i. \end{aligned}$$

Application of Lemma (ii) to the terms on the right of equation (11) yields

$$(12) \quad \sum_{k=1}^l a_k(t) X_k = \sum_{i=1}^l \sum_{k=1}^l g'_i(t) \xi_{ki} X_k.$$

Since the operators X_k are linearly independent we have a linear relation between the $a_k(t)$ and the $g'_i(t)$. The elements ξ_{ki} of the transform matrix ξ are analytic functions of the g_i 's.

$$(13) \quad \begin{matrix} a & \xi & g' \\ \begin{bmatrix} a_1 \\ \vdots \\ a_2 \\ \vdots \\ a_l \end{bmatrix} & = \begin{bmatrix} \xi_{11} & \cdots & \xi_{1l} \\ & & \\ \xi_{l1} & & \xi_{ll} \end{bmatrix} \begin{bmatrix} g'_1 \\ \vdots \\ g'_2 \\ \vdots \\ g'_l \end{bmatrix}, & g(0) = 0. \end{matrix}$$

Since the ξ_{ki} are analytic functions of g , we have that the determinant Δ of ξ is an analytic function of g . We also know that at $t=0$, $\xi=I$, and hence $\Delta(0) \neq 0$. These two facts show that there must exist a neighborhood N_0 of $t=0$ in which $\Delta \neq 0$, i.e., in which ξ is invertible. We can thus write (13) in the form

$$(14) \quad \frac{dg}{dt} = f(a, g) = \xi^{-1}a, \quad g(0) = 0, \quad t \in N_0.$$

Since ξ^{-1} is analytic in N_0 , we are assured of a neighborhood of $t=0$ in which the solution of (14) exists and is unique. This completes the proof of the theorem.

EXAMPLE. Let $dU/dt = (a_1(t)H + a_2(t)E + a_3(t)F)U$, where H, E , and F are constant linear operators with the Lie multiplication table $[E, F] = H; [E, H] = 2E; [F, H] = -2F$. Setting $U = e^{\theta_1(t)H}e^{\theta_2(t)E}e^{\theta_3(t)F}$, we obtain

$$\begin{aligned} a_1H + a_2E + a_3F &= g_1' H + g_2' (e^{\theta_1 adH})E + g_3' (e^{\theta_1 adH} e^{\theta_2 adE})F \\ (e^{\theta_1 adH})E &= E - 2g_1E + \frac{(2g_1)^2}{2!} E - \dots = e^{-2\theta_1}E \\ (e^{\theta_1 adH} e^{\theta_2 adE})F &= (e^{\theta_1 adH})(F + g_2H + g_2^2E) \\ &= e^{2\theta_1}F + g_2H + g_2^2 e^{-2\theta_1}E. \end{aligned}$$

Hence g_1, g_2, g_3 satisfy the equations

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 1 & 0 & g_2 \\ 0 & e^{-2\theta_1} & g_2^2 e^{-2\theta_1} \\ 0 & 0 & e^{2\theta_1} \end{bmatrix} \begin{bmatrix} g_1' \\ g_2' \\ g_3' \end{bmatrix}.$$

4. Global results.

THEOREM 2. *If \mathfrak{L} is solvable, then there exists a basis, and an ordering of this basis, for which Theorem 1 is global.*

PROOF. A theorem of Lie states that if \mathfrak{L} is solvable, then there exists a chain of ideals $0 \subset L_1 \subset L_{l-1} \subset \dots \subset L_l = \mathfrak{L}$, where each L_m is exactly of dimension $l - m + 1$. It is easy to see that there is a basis X_1, X_2, \dots, X_l for \mathfrak{L} which can be arranged so that L_m is the ideal generated by X_m, \dots, X_l . With this arrangement the multiplication table for \mathfrak{L} becomes

$$(15) \quad [X_i, X_j] = \sum_{k=i}^l \gamma_{ij}^k X_k \quad \text{for } i > j.$$

Since $(\prod_{j=1}^{i-1} \exp(g_j adX_j))X_i = \sum_{k=1}^i \xi_{ki} X_k = \sum_{k=i}^l \xi_{ki} X_k$, we have that $\xi_{ki} \equiv 0$ for $i > k$. It is clear that ξ_{ki} depends only on g_j with $j < k$; thus system $\xi g' = a$ is in triangular form. To show that $\xi g' = a$ has a solution for all t for which $a(t)$ is continuous, it is sufficient to prove that the diagonal elements ξ_{ii} never vanish. We shall show that $\xi_{ii} = \exp(\sum_{j=1}^{i-1} g_j \gamma_{ji}^i)$. This is readily seen since, by use of Lemma (i), one finds that

$$\begin{aligned} \exp(g_{i-1} adX_{i-1})X_i &= X_i + g_{i-1} \gamma_{i-1,i}^i X_i + \text{terms in } L_{i+1} \\ &\quad + (g_{i-1} \gamma_{i-1,i}^i)^2 / 2! X_i + \text{terms in } L_{i+1} + \dots \\ &= X_i \exp(g_{i-1} \gamma_{i-1,i}^i) + \text{terms in } L_{i+1}. \end{aligned}$$

The proof is completed by repetition of the above argument.

There is an interesting formula due to Zassenhaus [see 4, p. 661] which states that if X, Y generate the free algebra \mathfrak{L} , then $e^{X+Y} = e^X e^Y e^{C_2} e^{C_3} \cdots e^{C_n} \cdots$, where C_n is unique and exactly of degree n in X, Y . Magnus gives a method by which the C_n may be found recursively. For solvable algebras Theorem 2 yields the following sharpening of the Zassenhaus formula.

COROLLARY. *If X_1, \dots, X_m generate a solvable Lie algebra \mathfrak{L} of dimension l , then*

$$\exp\left(\sum_{i=1}^m a_i X_i\right) = \prod_{i=1}^l \exp(g_i Y_i),$$

where Y_1, \dots, Y_l is a suitable basis for \mathfrak{L} . The g_i can be found by quadrature.

Theorem 2 allows us to confine our attention to the case where \mathfrak{L} is semisimple. To see this we make use of Levi's theorem (see Jacobson [2, p. 91]); \mathfrak{L} may be decomposed into the direct sum $\mathfrak{L} = L_0 + L_1$, where L_0 is the radical of \mathfrak{L} , and L_1 is a semisimple subalgebra of \mathfrak{L} . In the equation $dU/dt = A(t)U$, where $A(t)$ generates \mathfrak{L} , the decomposition $\mathfrak{L} = L_0 + L_1$ gives rise to the corresponding decomposition, $A = A_0 + A_1$ of A , where A_i is in L_i . If we have proved that Theorem 1 is global for all semisimple Lie algebras as well as all solvable Lie algebras, then the following scheme extends the result to \mathfrak{L} . Let

$$(16) \quad \frac{dU}{dt} = AU = (A_0 + A_1)U.$$

Define U_0 , and U_1 , by $dU_1/dt = A_1 U_1$, $dU_0/dt = (U_1^{-1} A_0 U_1) U_0$. Since L_0 is an ideal in \mathfrak{L} , we see (by Lemma (i)) that $U_1^{-1} A_0 U_1$ is in L_0 . It is easy to verify that $U = U_1 U_0$ satisfies (16).

If \mathfrak{L} is semisimple, the global nature of Theorem 1 is in doubt. The examples below show that the choice of basis is critical.

EXAMPLE 1. Let $dU/dt = AU$ with A a 3×3 antisymmetric matrix. A generates the Lie algebra $X_1 = E_{12} - E_{21}$, $X_2 = E_{13} - E_{31}$, $X_3 = E_{23} - E_{32}$, $[X_1, X_2] = -X_3$, $[X_1, X_3] = X_2$, $[X_2, X_3] = -X_1$. If we assume a solution of the form $U = e^{g_1 X_1} e^{g_2 X_2} e^{g_3 X_3}$ and compute the corresponding ξ matrix, we find that $|\xi| = \cos g_2$. Hence ξ is not invertible for all time, and the representation is not global. We get the same result for any ordering of the above basis.

EXAMPLE 2. This example shows that even if ξ is invertible for all time, the representation by a product of exponentials need not be

global. Let $dU/dt = AU$ with A continuous for $0 \leq t < \infty$, $a_{11} = -a_{22}$, and

$$A = \begin{bmatrix} -6 & -4 \\ 5 & 6 \end{bmatrix} \quad \text{for } 0 \leq t < 1.$$

The Lie algebra generated by A is the same as that given in the example of §3. If we assume a solution of the form $U = \exp(g_1 H) \exp(g_2 E) \exp(g_3 F)$, we find that for $0 \leq t < 1$, $e^{g_1} = \cosh 4t - (3/2) \sinh 4t$. Thus, $g_1(t) = -\infty$ at $t = (1/8) \ln 5$, and the representation is not global. In this example $|\xi| \equiv 1$ so that ξ is always invertible.

The next example is interesting enough to state as a theorem.

THEOREM 3. *If $dU/dt = BU$, where B is any real continuous 2×2 matrix, then U has the form $U = \exp(g_1 K) \exp(g_2 A) \exp(g_3 N) \exp(g_4 I)$, where $K = E_{12} - E_{21}$, $A = E_{11} - E_{22}$, $N = E_{12}$, and I is the identity; this representation is global.*

PROOF. Let $B = S + aI$, S generates the algebra $\{K, A, N\}$. Then U has the form Ve^{aI} with $g_4 = \int_0^t a(\tau) d\tau$ and $dV/dt = SV$. Theorem 1 shows that V has the required form in a neighborhood of the origin. Now,

$$\begin{aligned} & \exp(g_1 K) \exp(g_2 A) \exp(g_3 N) \\ &= \begin{bmatrix} e^{g_2} \cos g_1 & g_3 e^{g_2} \cos g_1 + e^{-g_2} \sin g_1 \\ -e^{g_2} \sin g_1 & -g_3 e^{g_2} \sin g_1 + e^{-g_2} \cos g_1 \end{bmatrix} = \begin{bmatrix} V_{11} & V_{12} \\ V_{21} & V_{22} \end{bmatrix}. \end{aligned}$$

Solving for the g_i in terms of the V_{ij} gives $g_1 = -\text{Arctan}(V_{21}/V_{11})$, $g_2 = (1/2) \ln(V_{11}^2 + V_{21}^2)$, $g_3 = (V_{11}V_{12} + V_{21}V_{22})/(V_{11}^2 + V_{21}^2)$. Since V is nonsingular for all time, V_{11} and V_{21} cannot vanish simultaneously. Hence the g_i are analytic functions of the V_{ij} (V_{ij} assumed real). This completes the proof.

If $dU/dt = AU$, where A generates $\mathcal{L} = \{X_1, \dots, X_l\}$, then we may consider the representation $U = \prod \exp(g_i X_i)$ as a transformation of coordinates. In order that the above representation be global, it is clearly necessary and sufficient that (a) the mapping of the Lie algebra \mathcal{L} to the solution space of $dU/dt = AU$ is onto, and (b) the Jacobian, $J(U/g)$, of the transformation has rank l for all time. The second condition is fairly easy to investigate.

THEOREM 4. *Let $dU/dt = AU = (\sum_{i=1}^l a_i X_i)U$, and let ξ be the matrix of Theorem 1. Then the rank of J is the same as the rank of ξ whenever the two matrices are defined (i.e., whenever g_1, \dots, g_l are bounded).*

PROOF.

$$\frac{\partial U}{\partial g_k} = \prod_{i=1}^{k-1} \exp(g_i X_i) X_k \prod_{i=k}^l \exp(g_i X_i)$$

$$\frac{\partial U}{\partial g_k} U^{-1} = \left(\prod_{i=1}^{k-1} \exp(g_i \text{ad} X_i) \right) X_k = \sum_{i=1}^l \xi_{ik} X_i.$$

Therefore, $JU^{-1} = \xi X$, where X is the "vector" $(X_1, \dots, X_l)^T$. Since U^{-1} is nonsingular, J and ξ must have the same rank.

The next theorem shows that there do exist bases for which ξ is invertible.

THEOREM 5. *Let \mathfrak{L} be a split Lie algebra of dimension l . Then \mathfrak{L} can be decomposed, (e.g. the root space decomposition), into $\mathfrak{L} = L_1 \oplus L_2$ where L_1 and L_2 are solvable subalgebras. There exists a basis, and an ordering of this basis, for which the matrix ξ of Theorem 1 is always nonsingular.*

PROOF. For L_1 choose the basis and ordering as in Theorem 2, say X_1, \dots, X_m . Then ξ will have the form

$$\xi = \begin{bmatrix} A & * \\ 0 & B \end{bmatrix},$$

where A is upper triangular and nonsingular. Hence ξ is invertible if B is. Let Y_1, \dots, Y_n be a basis for B . The k th column of B consists of the coefficients of Y_1, \dots, Y_n in the expansion of $(\prod_{i=1}^m \exp(\text{ad} g_i X_i)) (\prod_{j=1}^{k-1} \exp(\text{ad} h_j Y_j)) Y_k$. Now, the operator $\prod_{i=1}^m \exp(\text{ad} g_i X_i)$ is an automorphism of \mathfrak{L} , so that it is enough to show that the matrix B_1 with k th column the coefficients of Y_1, \dots, Y_n in the expansion of $(\prod_{j=1}^{k-1} \exp(\text{ad} h_j X_j)) Y_k$ is nonsingular. B_1 is just the ξ -matrix for the algebra L_2 , and since L_2 is solvable, Y_1, \dots, Y_n can be chosen so that B_1 is nonsingular.

REFERENCES

1. N. Bourbaki, *Éléments de mathématique. Algèbres de Lie*, Actualités Sci. Ind. No. 26, Hermann, Paris, 1961.
2. N. Jacobson, *Lie algebras*, Interscience, New York, 1962.
3. J. Mariani and W. Magnus, *The exponential solution for the homogeneous linear differential equation of the second order*, Research report no. BE-37, New York University, 1961.
4. W. Magnus, *On the exponential solution of differential equations for a linear operator*, Comm. Pure Appl. Math. **7** (1954), 649-673.

SOCONY MOBIL OIL COMPANY, INC.